





- 1,284,900 new cancer cases will be diagnosed in the United States, including 14,800 in Iowa.
- 555,500 cancer deaths will occur in the United States, including 6,400 in Iowa.

The average annual age-adjusted mortality rate for cancer deaths per 100,000 persons²

lowa: 193.0 National: 206.0

Iowa ranks 40th highest overall in cancer mortality rates among the 50 states and Washington, D.C.

LUNG CANCER

In 2002, the American Cancer Society estimates¹

- 1,900 new cases of lung cancer will be diagnosed among men and women in Iowa.
- 1,700 men and women will die of lung cancer in Iowa.

The average annual age-adjusted mortality rates for lung cancer deaths per 100,000 persons, by race, 1995–1999³

	Iowa	National
Overall	52.9	57.7
White	52.8	57.5
Black	79.9	67.8
Hispanic	32.3	23.4
Asian/Pacific Islander	-	29.0
American Indian/Alaska Native	-	36.2

BREAST CANCER

In 2002, the American Cancer Society estimates¹

- 2,400 new cases of breast cancer will be diagnosed among women in Iowa.
- 500 women will die of breast cancer in Iowa.

The average annual age-adjusted mortality rates for breast cancer deaths per 100,000 women, by race, 1995–1999³

	Iowa	National
Overall	27.5	28.8
White	27.5	28.2
Black	34.4	37.1
Hispanic	-	17.2
Asian/Pacific Islander	-	13.0
American Indian/Alaska Native	-	15.0

COLORECTAL CANCER

In 2002, the American Cancer Society estimates¹

- 2,000 new cases of colorectal cancer will be diagnosed among men and women in Iowa.
- 800 men and women will die of colorectal cancer in Iowa.

The average annual age-adjusted mortality rates for colorectal cancer deaths per 100,000 persons, by race, 1995–1999³

	Iowa	National
Overall	22.9	21.7
White	22.9	21.3
Black	28.4	29.0
Hispanic	-	12.9
Asian/Pacific Islander	-	13.3
American Indian/Alaska Native	-	13.5

PROSTATE CANCER

In 2002, the American Cancer Society estimates¹

- 2,400 new cases of prostate cancer will be diagnosed among men in Iowa.
- 400 men will die of prostate cancer in Iowa.

The average annual age-adjusted mortality rates for prostate cancer deaths per 100,000 men, by race, 1995–1999³

	Iowa	National
Overall	33.1	33.9
White	32.9	31.2
Black	81.5	72.8
Hispanic	-	21.6
Asian/Pacific Islander	-	14.3
American Indian/Alaska Native	_	17.5

¹ Source: American Cancer Society Facts and Figures, 2002. Estimates exclude more than a million cases of basal and squamous cell skin cancers and in situ cancers, except urinary bladder, that will be diagnosed in 2002. Lung cancer rates include bronchus cancer. State death totals were rounded to nearest 100.

² Source: CDC National Center for Health Statistics, vital statistics data, underlying cause of death, 1995–1999. Death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population.

³ Hyphens represent suppression of rates when there were 75,000 or fewer persons in the denominator or 20 or fewer deaths in the numerator.